

Interesting Cases

Acquired Methemoglobinemia Secondary to Nitrobenzene Poisoning: A Case Report

Sumanth S Badhya¹, BK Srikanth²

¹Postgraduate, ²Associate Professor, Department of Paediatrics, Vijayanagar Institute of Medical Sciences, Ballari, Karnataka, India.

INFO

Corresponding Author:

Sumanth S Badhya, Department of Paediatrics, Vijayanagar Institute of Medical Sciences, Ballari, Karnataka, India.

E-mail Id:

bsumanthbadhya369@gmail.com

Orcid Id:

<https://orcid.org/0000-0001-5203-8455>

How to cite this article:

Badhya SS, Srikanth BK. Acquired Methemoglobinemia Secondary to Nitrobenzene Poisoning: A Case Report. Postgrad J Pediatr Adol Med. 2025;1(2):40-42.

Date of Submission: 2025-07-29

Date of Acceptance: 2025-12-17

ABSTRACT

Methemoglobinemia occurs as a result of the oxidation of iron in haemoglobin from the ferrous state to the ferric state. Acquired methemoglobinemia is one of the serious side effects occurring as a result of exposure to nitrate-containing compounds. Treatment of these patients usually involves the administration of methylene blue. Here, we have reported a 14-year-old adolescent with nitrobenzene poisoning.

Keywords: Poisoning, Methemoglobinemia, Nitrobenzene, Methylene Blue

Introduction

Adolescent poisoning is a common problem encountered in a paediatric emergency. Nitrobenzene is a potentially fatal compound used as a paint solvent and dye, and in rubber synthesis. Consumption of nitrobenzene leads to methemoglobinemia. Early diagnosis of methemoglobinemia and administration of methylene blue prevents mortality in these cases. We report a 14-year-old adolescent child with nitrobenzene poisoning.

Case Report

A 14-year-old child came to our paediatric emergency room with a history of an alleged suicidal attempt by consumption of 20% nitrobenzene (Figure 1) 9 hours prior to presentation. She had presented with pain in abdomen and vomiting since then. There was no history of headache, chest pain, hurried breathing, drowsiness or convulsions. On examination, she was conscious with a GCS of 14/15 with

an oxygen saturation of 60% in room air which improved to 68% with 5 L of oxygen. She had an odour of bitter almond with peripheral cyanosis. Except for mild epigastric tenderness, rest of the systemic examination was fairly normal.

A stomach wash was given. Blood was drawn, which appeared dark in colour (Figure 2) and was sent for investigations including blood gas analysis and serum methemoglobin levels. In view of hypoxia, a chest X-ray was done which was normal. Blood investigations showed raised bilirubin levels (max - 2 mg/dl), a slight drop in haemoglobin and serum methemoglobin levels of 6.2% (Figure 3). Rest of the investigations were normal. Injection methylene blue 1 mg/kg was given intravenously over 20 mins. This showed a dramatic improvement in her saturation to 86-88% within an hour. Two repeat doses of methylene blue were given 8 hours apart. She also received 500 mg/day of oral Vitamin C for 1 week. The following day her saturation improved to 92-94% at room air. Over the course of the



next few days, she showed good clinical improvement. On HEADSS screening, she was found to have a history of excessive usage of electronic gadgets (cell phones) and episodes of emotional outbursts and impulsive behaviour. Psychiatric assessment and counselling were done and she was discharged within a week.



Figure 1.20% Nitrobenzene Solution



Figure 2. Dark Brown Coloured Blood

Barcode	MAMATHA 279 V2558577 11208021926	Reference: Dr.AMRUTH DIAGNOSTIC LABORATORY BELLARY	VID: 11208021926
PID NO:	P11200434358	Sample Collected At: R v diagnostics laboratory	Registered On: 02/12/2020 02:22 PM
Age:	14 Year(s)	# 76/10 4th main 15 cross malleshwaram bangalore 560003 zone: out-01(os)	Collected On: 02/12/2020
Sex:	Female	PROCESSING LOCATION: Metropolis Healthcare Ltd, Unit No. 409-416, 4th Floor, Commercial Building-1, Kohinoor Mall, Mumbai-70	Reported On: 04/12/2020 03:15 PM

Investigation	Observed Value	Unit	Biological Reference Interval
Meth-haemoglobin** (Heparin Whole Blood Spectrometry)	6.2	%	< 1.5

Medical Remarks: Kindly correlate clinically.

Interpretation:

Methemoglobin accumulation in erythrocytes is due to acquired and hereditary causes.

- Acquired causes is most commonly encountered as a result of administration of medications such as local anaesthetic agents, especially procaine and benzocaine amyl nitrite. Other drugs like chloroquine, dapsone, nitrates, nitrites, nitrogllycerin, nitroprusside, phenacetin, phenazopyridine, primaquine, quinones and sulfonamides can also lead to accumulation of methemoglobin in RBCs as a result of drug intolerance or due to its toxicity.
- Congenital methemoglobinemias are rare.
- They are either due to:
 - A deficiency of methemoglobin reductase (also called cytochrome B5 reductase or diaphorase) in erythrocytes which is an autosomal recessive disorder
 - One of several intrinsic structural disorders of hemoglobin, called methemoglobin-M which are inherited as autosomal dominant.
 - G6PD deficiency
 - Pyruvate kinase deficiency

-- End of Report --

Figure 3. Serum Methemoglobin Levels
Discussion

Nitrobenzene is an important component in the synthesis of various solvents and flowering agents. It has a bitter almond odour and is yellowish in colour with an oily texture. The first case reporting poisoning due to nitrobenzene was seen in 1886.¹ Lethal dose of nitrobenzene is around 1 gm to 10 gm.²

It may be absorbed through the skin, respiratory tract, and gastrointestinal tract, and its highest concentration accumulates in the liver, brain, blood, and stomach due to its lipophilic nature.²

The ingestion of nitrobenzene results in a condition called methemoglobinemia. This condition causes the oxidation of iron present in the blood from the ferrous (Fe^{2+}) to the ferric (Fe^{3+}) state, which further leads to the blood turning brown, and the loss of the ability to transport oxygen.³ Physiologically, a low level (< 1%) of haemoglobin is oxidised to methemoglobin. This low level is maintained due to the presence of two reductive pathways in the red blood cells, namely diaphorase pathways and HMP

pathway. In nitrobenzene poisoning, due to oxidative stress, enhanced quantities of methemoglobin are produced.² There are various mechanisms helpful in the reduction of methemoglobin, for example, catalysis by cytochrome b5 reductase and alternate pathway using NADPH-dependent methemoglobin reductase system.⁴

In case of less than 10%-15% of methemoglobin, only cyanosis seems plausible (as in our case), however, the child will be usually asymptomatic. Levels above 20% cause tachypnoea, headache, chest pain, and dyspnoea. 40-50% of methemoglobin causes lethargy and confusion. It might also cause metabolic acidosis further resulting in seizures, coma, bradycardia, hypertension, and ventricular dysrhythmia. Methemoglobin levels above 70% are lethal. Some other effects are haemolytic anaemia, altered liver functions, contact dermatitis, and hepatosplenomegaly. Severe symptoms are generally found in children with anaemia and G6PD enzyme deficiency.⁵

Methylene blue is an effective antidote for methemoglobinaemia. The advised dosage is 1-2 mg/kg (with a maximum of 50 mg dose in adults) given intravenously as a 1% solution over 5 minutes. If required, it can be repeated in one hour. At levels above 7 mg/kg, methylene blue acts as an oxidant and therefore may cause methemoglobinaemia in some individuals. For patients with methemoglobin levels of more than 30%, ascorbic acid (200-500mg/day) may be advised. The use of N-acetylcysteine in the treatment of methemoglobinaemia has been reported. Patients showing severe symptoms may be advised exchange transfusion. Cases with methemoglobin levels > 50% or those who are not responding to standard treatment are provided hyperbaric oxygen.⁵

Conclusion

Nitrobenzene poisoning is rarely encountered in clinical practice. Hence high index of suspicion in cases of unknown compound consumption and the above-mentioned features should prompt us to think of acquired methemoglobinaemia as administration of methylene blue can be life saving in these cases.

Source of Funding: None

Conflict of Interest: None

References

1. Gupta G, Poddar B, Salaria M, Parmar V. Acute nitrobenzene poisoning. Indian Pediatr. 2000;37:1147-8. [PubMed] [Google Scholar]
2. Shrestha N, Karki B, Shrestha PS, Gami R, Acharya SP, Acharya S. Management of nitrobenzene poisoning with oral methylene blue and vitamin C in a resource limited setting a case report. Toxicol Rep. 2020;7:1008-9. [PubMed] [Google Scholar]
3. Lee CH, Kim SH, Kwon DH, Jang KH, Chung YH, Moon JD. Two cases of methemoglobinemia induced by the exposure to nitrobenzene and aniline. Ann Occup Environ Med. 2013;25(1):1-7. [PubMed] [Google Scholar]
4. Padyana M, Shetty AJ, Suresh PK. Nitrobenzene poisoning with methemoglobinemia. Indian J Case Rep. 2019;5(1):59-60. [Google Scholar]
5. Saxena H, Saxena AP. Acute methaemoglobinaemia due to ingestion of nitrobenzene (paint solvent). Indian J Anaesth. 2010;54(2):160. [PubMed] [Google Scholar]