

Review Article

Obesity Epidemiology: A Serious Public Health Concern in India

Bhanumati Sarkar¹, Paramita Biswas², Chandan Kumar Acharya³, Sandip Kumar Jana⁴,
Nasrin Nahar⁵, Souparna Ghosh⁵, Dipayan Dasgupta⁶, Sudipta Kumar Ghorai⁷,
Nithar Ranjan Madhu⁵

¹Department of Botany, Acharya Prafulla Chandra College, New Barrackpore, West Bengal, India.

²Department of Agronomy, Faculty of Agriculture, Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal, India.

³Department of Botany, Bajkul Milani Mahavidyalaya, Kismat Bajkul, West Bengal, India.

⁴Department of Zoology, Bajkul Milani Mahavidyalaya, Purba Medinipur, West Bengal, India.

⁵Department of Zoology, Acharya Prafulla Chandra College, New Barrackpore, West Bengal, India.

⁶Department of Computer Science and Engineering, BP Poddar Institute of Management and Technology, West Bengal, India.

⁷Department of Zoology, Egra SSB College, Purba Medinipur, West Bengal, India.

DOI: <https://doi.org/10.24321/2278.2044.202205>

I N F O

Corresponding Author:

Nithar Ranjan Madhu, Department of Zoology, Acharya Prafulla Chandra College, New Barrackpore, West Bengal, India.

E-mail Id:

nithar_1@yahoo.com

Orcid Id:

<https://orcid.org/0000-0003-4198-5048>

How to cite this article:

Sarkar B, Biswas P, Acharya CK, Jana SK, Nahar N, Ghosh S, Dasgupta D, Ghorai SK, Madhu NR. Obesity Epidemiology: A Serious Public Health Concern in India. Chettinad Health City Med J. 2022;11(1):21-28.

Date of Submission: 2022-01-11

Date of Acceptance: 2022-01-28

A B S T R A C T

Over the past several years, India has seen the incidence of obesity double or quadruple due to economic expansion, industrialisation, motorised transportation, urbanisation, a more sedentary lifestyle, and a nutritional change to prepare dishes that are rich in calories. Many things contribute to becoming overweight or obese since bodyweight is influenced by various factors, including genetics, metabolism, behaviour, the environment, culture, and socioeconomics. When evaluating the risk of various diseases, the quantity, location, and distribution of fat and obesity in the human body are significant. Being overweight and having an improper fat distribution are linked to a variety of illnesses. Obesity occurs when a person's weight exceeds their body fat percentage. Therefore, obesity has already increased dramatically in India as a consequence of these factors, putting Indian residents at risk for health concerns. Our study aims to bring awareness to the problems, better understand the problems, and record the incidence of obesity and overweight in the Indian community. Obesity education and public awareness may go a long way toward preventing these issues.

Keywords: Public Health, India, Nutritional Status, Obesity, Overweight

Introduction

Obesity is a physiological disease in which fat accumulates in the body. It's a chronic ailment that the WHO and numerous other national and international organisations have categorised as a disease.¹ The condition is increasingly

being recognised as a significant contributor to the global disease burden, owing to the pandemic levels of obesity that have been seen in recent decades. In industrialised countries, cancer and cardiovascular disease are the two main causes of mortality. Unfortunately, 'cardiovascular

Chettinad Health City Medical Journal (P-ISSN: 2277-8845 & E-ISSN: 2278-2044)

Copyright (c) 2022: Author(s). Published by Advanced Research Publications



disease (CVD) is a primary reason for death and disability. In heart disease survivors, the expense of 'blood pressure-lowering', 'antithrombotic', and 'diuretic' drugs is high. Stroke survivors are often plagued by mental health issues and physical problems such as weakness in their muscles and joints. Obesity's cost to society is borne almost entirely by diabetes.²⁻⁵

'Overweight' and 'obesity' have experienced a rise in global incidence in recent years. The WHO predicted that more than seven hundred million individuals and 2.3 billion people would be overweight by 2015. An estimated 2.5+ million Americans lose their lives each year as a result of obesity-related illnesses. Overconsumption of calories results in fat storage in the body. Obesity, caused by this fat storage in the body (Figure 1), is becoming one of the most frequent nutritional diseases in industrialised nations. As a result, one in every four persons will acquire diabetes. As a consequence of being overweight or obese, one in every three people will have 'heart disease', and one in every four people will acquire cancer.⁶⁻⁹




Figure 1. Complications of Obesity as a Public Health Problem (BioRender was used to make this image)

When someone's weight exceeds what is deemed normal or suitable for their height and weight, they are said to be overweight. When it comes to maintaining body mass (BMI), obesity is defined by a high amount of fat in the form of 'adipose tissue'.¹⁰⁻¹² Obesity is defined as having "too much fatness" or having obesity that causes illness.¹³ Even while it is impossible to identify the specific causes of the epidemic, a sedentary lifestyle, a lack of physical activity and an abundance of energy-dense meals are among the primary factors.¹⁴⁻¹⁸ As a result, obesity has risen to become the most serious worldwide issue confronting our nation.

The consequences of undernutrition and overnutrition for adolescents in West Bengal, India¹⁹⁻²³ and across India²⁴⁻²⁷ have been well documented. The most common reason

for someone to be undernourished is because they aren't eating enough. Individuals who suffer from this condition tend to lose bodyweight and have a set of symptoms, like sluggishness and osteoarthritis, among others (dry skin, low blood pressure etc.). In addition, studies have shown that prolonged hunger is linked to the development of nutritional oedema, as well as to burning feelings in the hands and feet.

System of Weight Classification for Adults

The weight-to-height ratio of the body is now the most commonly used criteria for obesity classification, and it varies from underweight or wastefulness (18.0+ kg/m²) to severe, even morbid, overweight (40.0 kg/m²). Measures of abdominal adiposity like waist circumference have become more important and discriminating in clinical and research settings as an overweight/ obesity predictor.²⁸ Adiposity in the abdominal region results from fat accumulation surrounding the organs, especially visceral fat. In addition, metabolic dysregulation, which is connected to an increased likelihood of cardiovascular disease and other ailments, is another effect of this fat.²⁹ European men must have a waist circumference less than 94 cm, and European women must have a waist circumference less than 80 cm if they want to reduce their cardiovascular risk. However, other races and ethnicities should use a different cutoff point.²⁹

BMI and Obesity Metrics Calculation

In the struggle against obesity, a standard public health measure of overweight and obesity should be established as soon as possible. The 'BMI' is a measure used by the 'National Institutes of Health' (NIH) to categorise people as overweight or obese. In order to calculate Body Mass Index, just two elements are required, height and weight, and their exact measurements are taken. Body Mass Index is obtained by dividing one's weight in kilogrammes by one's height in metres squared.³⁰

Possible Risks associated with Obesity

Calculating a result's 'attributable fraction' (the proportion of the outcome that can be attributed to obesity) is critical for determining whether or not obesity is contributing to an increased risk of disease, death, or disability.

$$AF_p = \frac{p(RR - 1)}{p(RR - 1) + 1}$$

The attributable fraction (AF), the proportion of men in each BMI range (p), and the related relative risk (RR) are all indicated by RR. Since the percentage of obese people in each attributable fraction calculation will be the same, relative risks may be compared to examine the varied effects of obesity on mortality and morbidity. This review was based on data gathered from a follow-up survey of health professionals.^{5,31-34} The nurses' health study in the United States was used to emphasise the relative risks

of “stroke”, “mortality”, “coronary heart disease”, and “type 2 diabetes mellitus” associated with different BMI categories.³⁵⁻³⁷

Microbiomes and Viruses as Environmental Pathogens

Despite the fact that numerous viruses have been identified as having the potential to have a causative role in obesity, higher Ad-36 viral loads have been seen. Studies undertaken in a range of human populations have shown that it is most likely the cause of obesity in both adults and children, with associations to other metabolic characteristics.^{38,39}

New insights regarding obesity and illnesses have come up as a result of recent discoveries in understanding how billions of bacteria in the human microbiome affect obesity, energy metabolism, and carbohydrate and fat digestion, for example, according to a research conducted on mice, greater adiposity is a feature that may be passed on by microbiome transplantation.^{40,41}

Diabetes

Type 2 diabetes is closely linked to diabetes and obesity, which should come as no surprise. A three-fold increased risk of type 2 diabetes is associated with being overweight, whereas a seven-fold increased risk is associated with being obese. Weight gain during early to mid-adulthood and obesity during childhood and adolescence are substantial risk factors for acquiring type 2 diabetes later in life (Table

1). When combined with other metabolic disorders (such as dyslipidemia, insulin resistance, hypertension, and poor glycaemic control), obesity doubles one’s chance of getting diabetes.^{42,43}

Cancer

It was discovered that obesity was a significant factor in around 6% of all cancers identified in 2007. Overweight people are more likely to develop gallbladder, oesophagus, colon, liver and pancreatic cancers; postmenopausal breast and ovarian cancer, endometriosis, and prostate and renal disease.^{44,45}

Heart and Vascular Diseases

Overweight and obesity are linked to their usual antecedents, hypertension and dyslipidemia in general, in addition to being well-known risk factors for ischemic stroke and heart disease. Recent research has repeatedly revealed that benign obesity seems to be a fiction contrary to popular belief. The presence of excess weight increases the chance of ‘heart disease’ and stroke and the consequences of hypertension, dyslipidemia, and hypoglycaemia. A recent study has focused on the link between childhood obesity and adult disease because of the high frequency of juvenile obesity. It has been seen that obesity during childhood or adolescence is connected with a two-fold or greater risk of “adult hypertension”, stroke, and CHD (coronary heart disease).^{46,47}

Table 1. Risk Factors, Comorbidities, and Consequences of Obesity

	Individual	Socioeconomic	Environmental
Risk factors (non-exhaustive)	Genetics	Lack of education	Junk foods (i.e., geographical locations with little or no quick access to healthful foods, for example, fresh produce and groceries) are a growing problem in America.
	Low levels of physical activity	Poverty	Obesogens are substances that cause weight gain (e.g., endocrine-disrupting chemicals)
	Calorie-dense foods that are deficient in nutritional value (e.g., sugar-sweetened beverages)		Resources for physical activity are scarce, and towns are not walkable.
	Excessive energy intake in comparison to energy requirements		Viruses
	Sleeping too little or too much		Microbiota
	Sedentariness		Obese social bonds are a problem

	Exposures during pregnancy and perinatal period		
	Drugs that are specific (e.g., steroids)		
	Conditions affecting one's psychological well-being (e.g., stress, depression)		
	Certain disorders (for example, Cushing's disease) are characterised by excessive production of cortisol.		
Comorbidities and Sequelae (non-exhaustive)	Hypertension	Type 2 diabetes	Dyslipidemia
	Heart and vascular illnesses are two of the most common types of cardiovascular disease.	Osteoarthritis	Infertility
	Cancers of a specific kind (e.g., colon, oesophageal, postmenopausal breast)	Liver illnesses are a group of disorders that affect the liver (e.g., non-alcoholic steatohepatitis, non-alcoholic fatty liver disease)	Conditions and illnesses of the respiratory system (e.g., asthma, sleep apnea)
	Gallstones	Treatment and survival after a traumatic event	Infection
	Disabilities due to physical limitations	Conditions affecting one's psychological well-being (e.g., psychosocial function, depression)	Absenteeism/ loss of productivity
	Years of life wasted due to premature death	Increased medical expenses	Absenteeism and decreased productivity

Obesity's Genetic Contribution

While genetics are likely to have a role, this very little variance in BMI, together with the large rise in obesity in both developed and developing nations over the last half-century, indicates additional obesity risk factors. An estimated 1.5% of the variance in BMI may be attributed to the 32 most prevalent genetic variations. The people with the highest genetic risk (those with an average BMI of 3.0 kg/m²) had more than 38 risk allele bearers (those with an average BMI of 2.7 kg/m²). An individual who is 5'3" (160 cm) tall has a 7 kg (11 lb) difference in weight between those with high and low hereditary risk.⁴⁸ The obesity pandemic cannot be explained by changes in the human population's genetic composition. It is possible that genes impact obesity development if individuals

react differently to settings that encourage inactivity and "high-calorie eating". Furthermore, genes instruct the organism on how to adapt to changes in the surrounding environment. Consequently, genetic variations that promote hunger and food intake may have a role in the occurrence of obesity. Occasionally, a single gene mutation may be identified as the cause of hereditary obesity in a family ("monogenic obesity"). The exact nature of the genetic and environmental interactions that lead to obesity, on the other hand, is still a mystery (multifactorial obesity). Identifying persons with a high risk of obesity-related diseases, including diabetes, heart disease, and certain malignancies, is common among healthcare professionals. The influence of shared genetics and environment may be seen in the health histories of close relatives. Families can't alter their DNA, but they can help their children develop

good behaviours, such as a nutritious diet and regular exercise. These changes might positively impact the health of current and future generations.^{49,50}

Obesity and Physical Training

Losing weight and being less obese may increase total energy expenditure and decrease adipose tissue mass. The ACSM guidelines include both aerobic and anaerobic exercises. Aerobic exercise uses up a significant amount of muscular oxygen (running, cycling, rowing, etc.). The amount of oxygen consumed during aerobic activity, on the other hand, meets the muscles' energy requirements without the need for additional energy.⁵¹

Insufficient oxygen is needed to fulfil the energy demands put on muscles during a weightlifting activity, making it an anaerobic activity. In order to generate more lactic acid and electricity, the muscles are required to break down other sources of energy.⁵²

Exercise includes "physical activity (PA)", even if it does not have to be part of an organised training programme or series of sessions. METS (Metric Equivalent Tons) are measures of energy consumption used to compare the amount of effort and energy consumed during metabolic equivalent activities to that expended when sitting passively. Regular exercise throughout the day and incorporating it into other facets of one's lifestyle are both emphasised in this piece.

Problem-solving, spare time physical activity, and transportation are all part of leading an active lifestyle. A person's respiratory, muscular and cardiovascular fitness are essential considerations. Several recent studies have shown the benefits of physical exercise for persons of all ages on all of these fronts.^{53,54}

Experts in West Bengal performed various studies on the impact of regular exercise on physiological health markers such as VO_2 Max, blood pressure, heart rate, and menstruation, and the association between menstruation and obesity.⁵⁵⁻⁶⁰

The teenage years are a period of fast growth and maturation. Teenage years are marked by increased dietary needs, which might vary substantially depending on your residence.⁶¹ Unfortunately, data on teenagers' physical and nutritional state is rare, especially among indigenous peoples. Consequently, a database with data from various sections of the nation needs to be built. Teens in West Bengal have been studied to see how their physical and nutritional development compares to teenagers in other states.

Discussion and Conclusion

It's difficult to fathom obesity. Thorough knowledge of how the epidemic's risk variables interact and feasible therapies for the epidemic's multi-levelled and complicated

underlying causes are urgently needed. There has also been a significant decrease in physical labour and transportation automation due to the rapid economic expansion in many regions of the globe over the previous century. Diabetes has grown from being an uncommon illness of the rich to becoming one of the most prevalent diseases, with a rising number of the poor suffering as a result of an increasingly sedentary lifestyle and widespread availability of calorie-dense foods, among other factors. Despite the fact that type 2 diabetes and cardiovascular disease are intimately connected, this rapid demographic change has significant ramifications. As a result, the time has come for us to focus on the problem at hand and take the necessary steps to fix it.

There are important clinical and epidemiological ramifications of obesity and other metabolic diseases that may be linked to disparities in body adiposity between men and women. There are persistent health issues connected with growing body mass index (BMI), and how BMI grading is interpreted in terms of risk differences depending on the community.

According to the study, this may be ascribed to these individuals being increasingly dependent on market economies. Consequently, their roles and responsibilities are changing from those associated with a subsistence economy toward those connected with wage labour and manufactured commodities. To summarise, India is now confronted with two separate nutrition concerns, with the potential for growing obesity-related public health burden in the future, while undernutrition remains a concern among children and adults.

Obesity was shown to be more common among poorer individuals in most research studies done throughout India. It was also shown to be more frequent among females than males in the state of New York. Following the outcomes of the research, it was discovered that the problem of overweight and obesity is more frequent in metropolitan regions. As a result, it is expected that an intensive preventative strategy would minimise the burden of many chronic comorbidities on India's healthcare system, such as diabetes, cardiovascular disease, hypertension (high blood pressure), infertility etc.

Overweight and obesity have become more common among adults and children in India in recent years. Most other emerging countries, including the United States and the United Kingdom, have seen similar trends. These forecasts may not account for regional and subnational differences due to the considerable diversity in cultures, cuisine, and economic development across all the states of India. Given that the Indian constitution devolves the responsibilities of administration of the state-level implementation of health and nutrition policies, as it does in the United States, studying how these estimates may shift at the state level

may be beneficial for the sake of future health policy planning. This might be accomplished by either establishing a distinct urban health programme or including a particular clause in the proposed legislation in National Urban Health Program that emphasises a healthy diet and frequent physical activity.

Conflict of Interest: None

References

1. Müller MJ, Geisler C. Defining obesity as a disease. *Eur J Clin Nutr.* 2017;71:1256-8. [Google Scholar]
2. World Health Organization. Obesity-preventing and managing the global epidemic. *Rep. WHO Consult. Obes. WHO/NUT/NCD/98.1.* WHO: Geneva, Switzerland; 1997.
3. Wolf AM, Colditz GA. Current estimates of the economic cost of obesity in the United States. *Obes Res.* 1998;6(2):97-106. [PubMed] [Google Scholar]
4. Rissanen A, Heliövaara M, Knekt P, Reunanen A, Aromaa A, Maatela J. Risk of disability and mortality due to overweight in a Finnish population. *BMJ.* 1990;301:835-7. [PubMed] [Google Scholar]
5. Visscher TL, Seidell JC. The public health impact of obesity. *Annu Rev Public Health.* 2001;22:355-75. [PubMed] [Google Scholar]
6. Thomas CS, Krishnaswami S. Distribution of Body Mass Index in Indian patients with coronary artery disease. *Indian Heart J.* 1995;47:134-7. [PubMed] [Google Scholar]
7. Bhadra M, Mukhopadhyay A, Bose K. Overweight and obesity among adult Bengalee Hindu women of Kolkata, India. *Human Ecology.* 2005;13(Spl Issue):77-83. [Google Scholar]
8. Roy CS, Mukhopadhyay A, Bhadra M. Prevalence of overweight and obesity among Bengalee urban adult men of North 24 Parganas, West Bengal, India. *Int J Exp Res Rev.* 2016;4:45-50. [Google Scholar]
9. Stunkard AJ, Wadden TA. *Obesity: theory and therapy.* 2nd ed. New York: Raven Press; 1993.
10. Khatun A, Bhadra M, Mukhopadhyay A, Bose K. Anthropometric assessment of nutritional status of Muslim adolescents of Deganga, North 24 Parganas, West Bengal, India. *Int J Exp Res Rev.* 2016;4:34-9.
11. Khatun A, Bhadra M, Mukhopadhyay A, Bose K. Nutritional status and effect of physical activity on anthropometric characteristics of Bengalee Muslim adolescents boys of North 24 Parganas, West Bengal, India. *Int J Exp Res Rev.* 2016;5:8-14.
12. Sarkar B, Ghorai SK, Jana SK, Dasgupta D, Acharya CK, Nahar N, Ghosh S, Madhu NR. Overweight and obesity in West Bengal: a serious public health issue. *VEETHIKA Int Interdisc Res J.* 2021;7(4):9-15. [Google Scholar]
13. Kopelman PG. Obesity as a medical problem. *Nature.* 2000;404:635-43. [PubMed] [Google Scholar]
14. Bhadra M, Paul P, Das T, Mukhopadhyay A. Physical growth pattern and nutritional status among adolescent Bhumij boys of Khatra Block, Bankura District, West Bengal, India. *Int J Exp Res Rev.* 2018;16:1-6. [Google Scholar]
15. Sinha R, Kapoor AK. Cultural practices and nutritional status among premenopausal women of urban setup in India. *Open Anthropol J.* 2010;3:168-71. [Google Scholar]
16. Roy CS, Mukhopadhyay A, Bhadra M. Age variations in obesity, adiposity and central body fat distribution among Bengalee urban adult male of North 24 Parganas, West Bengal, India. *Int J Exp Res Rev.* 2016;5:74-83.
17. Das P, Khatun A, Mukhopadhyay A, Bhadra M, Bose K. Nutritional status of adult Bengalee slum dwellers of Midnapore town, Paschim Medinipore, West Bengal, India. *Int J Exp Res Rev.* 2016;8:23-8.
18. Das P, Khatun A, Bhadra M, Mukhopadhyay A, Bose K. Anthropometric characteristics of adult Bengalee slum dwellers of Midnapore town, Paschim Medinipore, West Bengal, India. *Int J Exp Res Rev.* 2016;8:1-8.
19. Sarkar S. Livelihood strategies of street children using the urban space: a case study at Sealdah station area, Kolkata. *Int J Exp Res Rev.* 2016;7:44-52.
20. Mistri A. Nutritional status and haemoglobin level among adult Bengalee women in a sub-urban area in West Bengal. *Int J Exp Res Rev.* 2016;8:81-91.
21. Mitra M, Mukhopadhyay A, Bhadra M. Sex variations in anthropometric variables of Santal children of Birbhum district, West Bengal, India. *Int J Exp Res Rev.* 2017;10:30-6. [Google Scholar]
22. Bhadra M, Mitra M, Baul S, Mukhopadhyay A. Assessment of undernutrition among Santal children of Bolpur-Sriniketan block of Birbhum District, West Bengal, India. *Int J Exp Res Rev.* 2018;15:9-15. [Google Scholar]
23. Chakraborty D, Ghosh PN. Impact of backwardness on health-case study Pakhiralaya village, Gosaba Block, Sundarban, West Bengal, India *Int J Exp Res Rev.* 2019;20:28-39.
24. Maiti M. Low birth weight is associated with maternal nutrition of Indian women. *Int J Exp Res Rev.* 2017;12:24-30. [Google Scholar]
25. Devi U, Mahanta B, Borah PK, Das JK, Barman N, Rabha J, Borah A, Mahanta J. Menstrual hygiene practices among adolescent girls in rural areas of Dibrugarh: an exploration into the need for health promotion activity. *Int J Exp Res Rev.* 2017;13:1-9.
26. Algur K, Gawari A, Mohan K. Demographic inequality among the tribal and non-tribal community in Nasik district of Maharashtra State. *Int J Exp Res Rev.* 2017;13:10-17.

27. Madhu NR, Sarkar S. Present status of dietary fat and obesity. UGC-Sponsored National Seminar on Food security and sustainable nutrition in India: The present scenario. Acharya Prafulla Chandra College, New Barrackpore; 2016. p. 120-4.
28. Hu FB. Obesity and mortality: watch your waist, not just your weight. *Arch Intern Med.* 2007;167(9):875. [PubMed] [Google Scholar]
29. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, Fruchart JC, James WP, Loria CM, Smith Jr SC; International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; International Association for the Study of Obesity. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society and International Association for the Study of Obesity. *Circulation.* 2009;120(16):1640-5. [PubMed] [Google Scholar]
30. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: executive summary. Expert Panel on the Identification, Evaluation, and Treatment of Overweight in Adults. *Am J Clin Nutr.* 1998;68(4):899-917. [PubMed]
31. Baik I, Ascherio A, Rimm EB, Giovannucci E, Spiegelman D, Stampfer MJ, Willett WC. Adiposity and mortality in men. *Am J Epidemiol.* 2000;152:264-71. [PubMed] [Google Scholar]
32. Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes Care.* 1994;17:961-9. [PubMed] [Google Scholar]
33. Rimm EB, Stampfer MJ, Giovannucci E, Ascherio A, Spiegelman D, Colditz GA, Willett WC. Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol.* 1995;141:1117-27. [PubMed] [Google Scholar]
34. Walker SP, Rimm EB, Ascherio A, Kawachi I, Stampfer MJ, Willett WC. Body size and fat distribution as predictors of stroke among US men. *Am J Epidemiol.* 1996;144:1143-50. [PubMed] [Google Scholar]
35. Carey VJ, Walters EE, Colditz GA, Solomon CG, Willett WC, Rosner BA, Speizer FE, Manson JE. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. *Am J Epidemiol.* 1997;145:614-9. [PubMed] [Google Scholar]
36. Manson JE, Colditz GA, Stampfer MJ, Willett WC, Rosner B, Monson RR, Speizer FE, Hennekens CH. A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med.* 1990;322:882-9. [PubMed] [Google Scholar]
37. Rexrode KM, Hennekens CH, Willett WC, Colditz GA, Stampfer MJ, Rich-Edwards JW, Speizer FE, Manson JE. A prospective study of body mass index, weight change, and risk of stroke in women. *JAMA.* 1997;277:1539-45. [PubMed] [Google Scholar]
38. McAllister EJ, Dhurandhar NV, Keith SW, Aronne LJ, Barger J, Baskin M, Benca RM, Biggio J, Boggiano MJ, Eisenmann JC, Elobeid M, Fontaine KR, Gluckman P, Hanlon EC, Katzmarzyk P, Pietrobelli A, Redden DT, Ruden DM, Wang C, Waterland RA, Wright SM, Allison DB. Ten putative contributors to the obesity epidemic. *Crit Rev Food Sci Nutr.* 2009;49(10):868-913. [PubMed] [Google Scholar]
39. Parra-Rojas I, Moral-Hernandez OD, Salgado-Bernabé AB, Guzmán-Guzmán IP, Salgado-Goytia L, Muñoz-Valle JF. Adenovirus-36 seropositivity and its relation with obesity and metabolic profile in children. *Int J Endocrinol.* 2013;2013:463194. [PubMed] [Google Scholar]
40. Atkinson RL, Dhurandhar NV, Allison DB, Bowen RL, Israel BA, Albu JB, Augustus AS. Human adenovirus-36 is associated with increased body weight and paradoxical reduction of serum lipids. *Int J Obes (Lond).* 2005;29(3):281-6. [PubMed] [Google Scholar]
41. Almgren M, Atkinson R, He J, Hilding A, Hagman E, Wolk A, Thorell A, Marcus C, Näslund E, Östenson CG, Schalling M, Lavebratt C. Adenovirus-36 is associated with obesity in children and adults in Sweden as determined by rapid ELISA. *PLoS One.* 2012;7(7):e41652. [PubMed] [Google Scholar]
42. Clarke SF, Murphy EF, Nilaweera K, Ross PR, Shanahan F, O'Toole PW, Cotter PW. The gut microbiota and its relationship to diet and obesity: new insights. *Gut Microbes.* 2012;3(3):186-202. [PubMed] [Google Scholar]
43. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature.* 2006;444(7122):1027-31. [PubMed] [Google Scholar]
44. Abdullah A, Peeters A, de Courten M, Stoelwinder J. The magnitude of association between overweight and obesity and the risk of diabetes: a meta-analysis of prospective cohort studies. *Diabetes Res Clin Pract.* 2010;89(3):309-19. [PubMed] [Google Scholar]
45. Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes (Lond).* 2011;35(7):891-8. [PubMed] [Google Scholar]

46. Polednak AP. Estimating the number of U.S. incident cancers attributable to obesity and the impact on temporal trends in incidence rates for obesity-related cancers. *Cancer Detect Prev.* 2008;32(3):190-9. [PubMed] [Google Scholar]
47. Vainio H, Bianchini F. *Weight control and physical activity.* IARC Press; 2002. [Google Scholar]
48. Flint AJ, Hu FB, Glynn RJ, Caspard H, Manson JE, Willett WC, Rimm EB. Excess weight and the risk of incident coronary heart disease among men and women. *Obesity.* 2010;18(2):377-83. [PubMed] [Google Scholar]
49. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, Allen HL, Lindgren CM, Luan J, Mägi R, Randall JC, Vedantam S, Winkler TW, Qi L, Workalemahu T, Heid IM, Steinthorsdottir V, Stringham HM, Weedon MN, Wheeler E, Wood AR, Ferreira T, Weyant RJ, Segrè AV, Estrada K, Liang L, Nemesh J, Park JH, Gustafsson S, Kilpeläinen TO, Yang J, Bouatia-Naji N, Esko T, Feitosa MF, Kutalik Z, Mangino M, Raychaudhuri S, Scherag A, Smith AV, Welch R, Zhao JH, Aben KK, Absher DM, Amin N, Dixon AL, Fisher E, Glazer NL, Goddard ME, Heard-Costa NL, Loos RJ. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet.* 2010;42(11):937-48. [PubMed] [Google Scholar]
50. Bouchard C. Defining the genetic architecture of the predisposition to obesity: a challenging but not insurmountable task. *Am J Clin Nutr.* 2010;91:5-6. [PubMed] [Google Scholar]
51. Choquet H, Meyre D. Genetics of obesity: what have we learned? *Curr Genomics.* 2011;12:169-79. [PubMed] [Google Scholar]
52. Bateman LA, Slentz CA, Willis LH, Shields AT, Piner LW, Bales CW, Houmard JA, Kraus WE. Comparison of aerobic versus resistance exercise training effects on metabolic syndrome (from the Studies of a Targeted Risk Reduction Intervention Through Defined Exercise-STRRIDE-AT/RT). *Am J Cardiol.* 2011;108(6):838-44. [PubMed] [Google Scholar]
53. Bhadra M, Baul S, Mahapatra B, Mukhopadhyay A. Evaluation of health and nutritional status of adolescent Muslim of North Dum Dum, West Bengal, India. *Int J Exp Res Rev.* 2017;11:66-72.
54. Bechara RG, Kelly ÁM. Exercise improves object recognition memory and induces BDNF expression and cell proliferation in cognitively enriched rats. *Behav Brain Res.* 2013; 245:96-100. [PubMed] [Google Scholar]
55. Basak S. The relationship between core muscle stability and balance in yoga. *Int J Exp Res Rev.* 2019;19:49-52.
56. Basak S, Biswas K. A study of selective physiological parameters in physical training college students. *Int J Exp Res Rev.* 2016;3 1-6.
57. Basak S, Dutta SA. Comparative study of physical fitness parameters between general college students and training college students. *Int J Exp Res Rev.* 2016;4:26-30.
58. Basak S, Hansda K. Relationship of state anxiety and trait anxiety between physical education students and general degree college students. *Int J Exp Res Rev.* 2016;5:15-8.
59. Mandal S. The nutritional health factors of cashewnut (*Anacardium occidentale*, L.). *Int J Exp Res Rev.* 2016;7:18-20.
60. Pramanik B. A comparative study on the knowledge, attitude and risk perception regarding complications of type-2 diabetes mellitus between male and female diabetic patients attending diabetic clinics in selected hospital of West Bengal, India. *Int J Exp Res Rev.* 2018;15:16-27. [Google Scholar]
61. World Health Organization. *Physical Status: the use and interpretation of anthropometry.* Technical Report Series No. 854. World Health Organization, Geneva; 1995.