

Research Article

Workplace Safety and Security in Indian Healthcare Settings: A Cross-Sectional Survey

Kartik Chadhar¹, Richa Mishra², Semanti Das³, Indra Shekhar Prasad⁴, Prakalp Gupta⁵

¹Department of Community Medicine, Vardhaman Mahavir Medical College and Safdarjung Hospital, New Delhi, India.

²Department of Paediatrics and Preventive Dentistry, Centre for Dental Education and Research, AIIMS, New Delhi, India.

³Centre for Community Medicine, All India Institute of Medical Sciences, New Delhi, India.

⁴Department of Medicine, All India Institute of Medical Sciences, New Delhi, India.

⁵Centre for Community Medicine, All India Institute of Medical Sciences, New Delhi, India.

DOI: <https://doi.org/10.24321/2455.7048.202403>

I N F O

Corresponding Author:

Kartik Chadhar, Department of Community Medicine, Vardhaman Mahavir Medical College and Safdarjung Hospital, New Delhi, India.

E-mail Id:

kartikchadhar@gmail.com

Orcid Id:

<https://orcid.org/0009-0000-1063-9500>

How to cite this article:

Chadhar K, Mishra R, Das S, Prasad I S, Gupta P. Workplace Safety and Security in Indian Healthcare Settings: A Cross-Sectional Survey. *Epidem Int.* 2024;9(3):1-12.

Date of Submission: 2024-09-16

Date of Acceptance: 2024-09-22

A B S T R A C T

Introduction: Workplace safety and security are critical concerns in healthcare settings, where incidents of violence against healthcare workers are increasingly reported. This study aimed to assess the current status of security and safety measures in Indian healthcare institutions, focusing on the perceptions of healthcare workers across various settings.

Method: A cross-sectional survey was conducted among 1,566 healthcare workers from diverse medical institutions across India using a pre-tested, self-administered online questionnaire, which assessed various dimensions of workplace safety. Logistic regression was used to find differences among groups. Open-ended responses were analysed using thematic analysis.

Results: The findings revealed that 58.2% of healthcare professionals felt unsafe at the workplace due to various threats. Perception of low safety was more in undergraduates (74.2%) than faculty (51%). A high proportion of health workers (78.4%) reported having been threatened on duty. Nearly half of the healthcare professionals lacked a dedicated duty room. The existing duty rooms were found to be grossly inadequate in providing the basic amenities. State-funded public healthcare settings were found to have poor living conditions, inadequate security measures, and a low perception of safety as compared to private institutions. Male healthcare professionals had higher odds of feeling unsafe at the workplace.

Conclusion: The study highlighted the significant gaps in the security infrastructure within Indian healthcare settings. Over half of the health workforce feels that their workplace is unsafe, particularly in state and central government medical colleges. A comprehensive, need-based approach with legislative support is necessary to enhance workplace security, which may improve doctor-patient relationships, workforce well-being, and overall healthcare quality.

Keywords: Healthcare Safety, Workplace Security, Violence at Workplace, Occupational Health, Security Measures

Introduction

Safety and security are essential processes to ensure the right to live and any form of threat to the same is a violation of fundamental human rights.¹ For the growth and development of human society, which depends on healthcare workers, effective and quality healthcare services are required. The health workforce includes those that provide health services such as doctors, nurses, etc, and those that support health services such as hospital managers, ambulance drivers, etc.² The lack of proper security measures renders the healthcare workers to work in unsafe environments with the perception of fear. Such circumstances further make the overall environment in the healthcare setup vulnerable to security threats and violence. Such acts of violence against healthcare workers have been repeatedly reported in literature.³ Along with healthcare professionals, property, equipment, and records have also been reported to be at risk of damage. Healthcare professionals, under these environments, may feel threatened to work, remain mentally burnt out, and may have high levels of stress, and therefore may not be able to provide patient care to their fullest potential.⁴ These safety and security issues are some of the causes of reluctance for most of the young doctors to go to rural health centres.⁵

One of the primary causes of such an incidence of violence may be attributed to a lack of safety and security in healthcare setups. A thorough literature search revealed a dearth of information about the safety and security in these setups, especially in the Indian subcontinent. A security lax, often in association with long duty hours of healthcare workers in substandard quality working environments, inadequately staffed hospitals, and poor health literacy among the general population may further aggravate the problem.^{3,6} This becomes particularly important in the case of students and residents who are usually found to be burdened with hectic duty hours, apart from rigorous academic schedules. The dissatisfaction among patients associated with a lack of adequate equipment, drugs, or other materials and sometimes a long waiting list for certain procedures, is often dealt with by these doctors. Furthermore, these doctors are also closely associated with the families during vulnerable times, such as patient mortality, which may lead to emotional outbursts among relatives.⁷⁻⁹

Due to the lack of an institutional mechanism to conduct regular surveys or audits of security services, the demand for security measures becomes incongruent with the security services in place. This is especially relevant in the current times when workplace violence against doctors is on the rise¹⁰ and the importance of workplace security is being highlighted. The National Task Force of the Union

Health Ministry has also invited recommendations for the same.¹¹ The current study shall help in the thorough understanding of the status of security and safety measures and the critical points for action. Therefore, this study aims to assess the security and safety measures in place in healthcare settings in India. The findings will also help in framing guidelines to strengthen security as well as serve as a baseline for estimating progress toward achieving a safe and secure workplace.

Materials and Method

We conducted a cross-sectional survey among healthcare workers all over India. A pre-tested, self-administered questionnaire was used to collect data on the status of security and safety services in their health institutions. To conduct the survey, an online form was circulated among healthcare workers (HCWs) from all cadres and disciplines across the country. Since obtaining a sampling frame was nearly impossible, we used convenient sampling to reach an adequate sample size. The calculated sample size was estimated to be 1688, assuming that the proportion of HCWs who felt unsafe at work was 50%, adjusting for a 10% non-response rate at 95% confidence levels and 5% relative precision. Data was collected in August 2024 through Google Forms shared across messaging platforms, e-mails, and social media.

The questionnaire was designed as a study tool to assess the current state of security and safety measures within healthcare settings. The survey gathered responses across several key areas, starting with demographic information. This section included questions about the respondent's location, age, gender, specialisation, designation, and the type of institution they were affiliated with, such as a government hospital or a private setup.

The survey then explored the perception of safety among respondents, focusing on how safe they felt within the hospital environment. It covered specific aspects like safety during night shifts and personal experiences with various threats, including physical, verbal, and sexual threats. A section on Security Infrastructure and Measures evaluated the adequacy of security measures such as the presence of security personnel, surveillance cameras, access control systems, emergency alarms, and lighting conditions. It also assessed the status of facilities like duty rooms, cleanliness, ventilation, and availability of basic amenities (e.g., washrooms). The Incident Reporting and Emergency Response section investigated whether respondents had witnessed violence, their awareness of emergency protocols, and the effectiveness of existing incident-handling procedures. A few open-ended questions explored in detail the nature and types of threats faced by healthcare workers and received suggestions to improve the current status of safety and security at the workplace.

In this article, the qualitative responses are presented as a summary. A detailed report on the qualitative responses will be presented in a follow-up article.

Data was analysed in R software after cleaning it in Microsoft Excel. The socio-demographic data was reported in percentages and frequencies. Missing data was addressed using multiple imputations. Continuous data such as age was checked for normality using the Shapiro-Wilk test. Normally distributed variables were reported using mean and standard deviation and non-normally distributed variables were reported with median and inter-quartile range. Other variables were reported in frequencies and percentages. Bivariate analysis was performed to find the association between the independent variables and dependent variables such as the perception of safety at the workplace based on gender, designation, department, and type of healthcare setting. All the variables having a significance level of ≤ 0.25 were included in the multivariate logistic regression model. Adjusted odds ratios (aOR) were reported for the predictors of dependent variables. Open-ended questions were analysed to identify the key themes using thematic analysis.

We obtained online informed consent from all the respondents after providing them the purpose of the survey.

Since we did not mandatorily collect the identity of the respondents, participation was completely voluntary, and data was presented as aggregates and percentages.

Results

Respondents' Characteristics

A total of 1627 responses were received from across India, of which, 1566 (96.2%) responses were found to be complete. Females constituted 55.5% of the total respondents. The age of the respondents ranged from 19 years to 81 years with the mean age being 30.9 ± 8.7 years. Almost one-fourth of responses (24.7%) were received from healthcare workers in Delhi. Almost half of the respondents were resident doctors (49.6%), followed by undergraduate medical students including interns (15.9%). Responses were also received from faculty members, medical officers, nursing staff, and other supporting staff. The majority of the respondents worked in government-run medical colleges (71.5%). Half of the respondents worked in non-surgical departments (49.2%) and one-third worked in surgical departments (33.8%). Other respondents included nursing, administrative, and other departments as detailed in Table 1.

Table 1. Characteristics of Respondents

Demographic Variable	Sub-Category	n (%)
Gender (N = 1566)	Female	869 (55.5)
	Male	697 (44.5)
Type of institution (N = 1560)	Central government medical college and hospital	394 (25.3)
	State government medical college and hospital	720 (46.2)
	Private medical college and hospital	157 (10.1)
	District hospital	78 (4.9)
	Community health centre	39 (2.5)
	Primary health centre	87 (5.6)
	Private clinic	35 (2.2)
	Others	50 (3.2)
Designation (N = 1562)	Faculty/ consultant	168 (10.8)
	Resident doctors	775 (49.6)
	Undergraduates and interns	249 (15.9)
	Medical officers	151 (9.7)
	Nursing staff	201 (12.8)
	Other supporting staff	18 (1.2)
Department (N = 1223)	Surgical departments	414 (33.8)
	Non-surgical departments	601 (49.2)
	Diagnostic and laboratory	99 (8.1)
	Nursing	78 (6.4)
	Administrative	5 (0.4)
	Others	26 (2.1)

Perception of Safety among Healthcare Professionals

Out of the total respondents, 624 (39.7%, 95% Confidence Interval (CI): 37.3–42.1) perceived the environment as unsafe and 291 (18.5%, 95% CI: 16.6–20.5) perceived it as very unsafe, indicating that more than half (915, 58.2%, 95% CI: 55.8–60.6) of the healthcare workers felt unsafe at the workplace.

Among the respondents, 1308 (83.4%) had worked a night shift, and of those, 821 (62.8%) reported feeling unsafe or very unsafe on the hospital premises during the night shift and only 166 (12.7%) felt safe. Almost four-fifths of the respondents (1222, 78.4%) reported having been threatened on duty, with the majority of threats being verbal (64%), 12% being physical, and about one-fourth (23.1%) being physical, verbal, and sexual in nature. On comparing the type of threat received among genders, it was observed that males had significantly higher odds (OR: 4.2, 95% CI: 2.92–6.03, $p < 0.01$) of reporting physical threats compared to females. On the other hand, females were 54% more likely than males to report experiencing multiple types of threats; verbal, physical and sexual. Both genders reported verbal threats at similar rates. Females were 8 times more likely (OR = 8.1, 95% CI: 1.0–63.4, $p = 0.046$) to report sexual threats than males although the precision was very low as only a few respondents reported sexual threats.

Determinants of Perceived Overall Safety and Night-Time Insecurity in Healthcare Settings

It was found that males had higher odds of perceiving themselves as unsafe in the workplace (aOR: 1.28, 95% CI: 1.02–1.61, $p = 0.031$) as compared to females. As compared to faculty members or consultants (168, 10.8%), resident doctors (775, 49.5%) showed an aOR of 1.22 (95% CI:

0.83–1.8, $p = 0.315$), and undergraduates/ interns (249, 15.9%) had an aOR of 1.48 (95% CI: 0.92–2.37, $p = 0.102$), with neither group reaching statistical significance. The perception of safety did not significantly vary across medical officers, nursing staff, and other supporting staff as well. These findings suggest a variation in perceived safety by gender but not significantly by professional designation. Compared to private medical colleges and hospitals, healthcare workers of state government medical colleges and hospitals had a significantly higher perception of unsafety (aOR: 6.44, 95% CI: 4.36–9.5, $p < 0.001$), similar to those of district hospitals (aOR: 4.15, 95% CI: 2.28–7.55, $p < 0.001$), community health centres (aOR: 6.09, 95% CI: 2.66–13.91, $p < 0.001$), and primary health centres (aOR: 5.14, 95% CI: 2.62–10.06, $p < 0.001$). Central government medical colleges and hospitals also showed increased unsafety (aOR: 1.55, 95% CI: 1.04–2.33, $p = 0.033$), while private clinics (aOR: 1.67, 95% CI: 0.75–3.7, $p = 0.209$) and ‘Other’ settings (aOR: 1.73, 95% CI: 0.87–3.44, $p = 0.115$) did not differ significantly. These findings indicate higher perceived unsafety in public healthcare settings compared to private ones as indicated in Table 2.

On analysing the perception of night security, females reported a slightly higher perception of night-time insecurity compared to males, but the difference was not statistically significant (aOR: 1.24, 95% CI: 0.98–1.57, $p = 0.069$). Regarding designation, no statistically significant differences were observed among various professional designations when compared to faculty members or consultants. Compared to private medical colleges and hospitals, all public healthcare settings showed a significantly higher perception of night-time insecurity, with the highest levels of insecurity observed in state government medical colleges (aOR: 5.78, $p < 0.001$). Private clinics also had increased insecurity (aOR: 2.49, $p = 0.026$), while ‘Other’ settings

Table 2. Perception regarding Safety According to Characteristics of Healthcare Professionals

Category	Sub-Category	Perception of Overall Safety		Total N (%)	Adjusted Odds Ratio (95% CI)	p Value
		Safe n (%)	Unsafe n (%)			
Gender	Female	394 (45.4)	475 (54.6)	869 (55.5)	Reference	-
	Male	259 (37.1)	438 (62.4)	697 (44.5)	1.28 (1.02–1.61)	0.031
Designation	Faculty/ consultant	89 (52.9)	79 (47.0)	168 (10.8)	Reference	-
	Resident doctors	339 (43.7)	436 (56.2)	775 (49.5)	1.22 (0.83–1.8)	0.315
	Undergraduates and interns	69 (27.7)	180 (72.2)	249 (15.9)	1.48 (0.92–2.37)	0.102
	Medical officers	59 (39.0)	92 (60.9)	151 (9.7)	1.00 (0.56–1.77)	0.994
	Nursing staff	88 (43.7)	113 (56.2)	201 (12.9)	1.02 (0.63–1.63)	0.943
	Other supporting staff	9 (52.9)	8 (47.1)	18 (1.2)	0.96 (0.33–2.79)	0.933

Type of healthcare setting	Number (n, %)	Unsafe (n, %)	Very Unsafe (n, %)	Reference	p-value
Private medical college	110 (70.1)	47 (29.9)	157 (10.1)	Reference	-
Central government medical college	236 (59.9)	158 (40.1)	394 (25.2)	1.55 (1.04–2.33)	0.033
State government medical college	182 (25.3)	538 (74.7)	720 (46.2)	6.44 (4.36–9.5)	< 0.001
District hospital	29 (37.7)	48 (62.3)	78 (5.0)	4.15 (2.28–7.55)	< 0.001
Community health centre	12 (30.8)	27 (69.2)	39 (2.5)	6.09 (2.66–13.91)	< 0.001
Primary health centre	30 (34.5)	57 (65.5)	87 (5.6)	5.14 (2.62–10.06)	< 0.001
Private clinic	22 (62.9)	13 (37.1)	35 (2.2)	1.67 (0.75–3.7)	0.209
Others	30 (60.0)	20 (40.0)	50 (3.2)	1.73 (0.87–3.44)	0.115

showed no significant difference.

It was seen that healthcare workers involved with state government healthcare institutions felt more unsafe (74.7%, 538/720) compared to the ones in central government healthcare institutions (40%) and a greater number of healthcare professionals working in government institutions felt more unsafe compared to individuals working in private healthcare institutions (30%) (Figure 1). Among the participants, 185 undergraduates (74.2%), 58 interns (68%), 91 medical officers (62%), and 346 junior residents (60%)




Figure 1. Perception regarding Safety Based on the Type of Health Institution

State government medical college (SGMC): Very safe - 1%, Central government medical college (CGMC): Very safe - 2%, Private medical college: Very safe - 5%




Figure 2. Perception regarding Safety Based on the Designation of Healthcare Professionals

felt the most unsafe with regard to the safety perception across various healthcare institutions (Figure 2).

Living Conditions of Healthcare Professionals During Duty Hours

As per the respondents, 781 (49.9%) healthcare professionals did not have a dedicated duty room to rest within the working area. If the duty room was present, it was not equipped with a washroom in 62.8% of the cases. Of the total respondents, 78.1% reported the lack of a separate duty room for female healthcare professionals, which included female doctors, nurses, and other female medical staff, and 69.5% reported the lack of a separate washroom for female health professionals. A staggering 1288 (82.3%) respondents reported about the inadequate number of beds in the duty rooms.

Duty Room Condition

The condition of the duty rooms was reported to be bad with inadequate cleanliness (68.8%), inadequate




Figure 3. Perception regarding Adequacy of Duty Room Essentials

room space (73.3%), pest infestation (81.1%), inadequate ventilation (70.6%), and lack of adequate air conditioning (64.2%) being the major problems reported (Figure 3).

Variations in the Condition of Duty Rooms among Medical Colleges in India

The adequacy of duty room amenities varied significantly across different types of medical colleges. State government medical colleges consistently showed higher odds of inadequacy compared to private medical colleges,

particularly for air conditioning (77% inadequate; OR: 2.12, $p < 0.01$), room space (80.5% inadequate; OR: 2.43, $p < 0.01$), ventilation (77.3% inadequate; OR: 1.88, $p = 0.01$), cleanliness (79.5% inadequate; OR: 2.98, $p < 0.01$), pest control (90.6% inadequate; OR: 4.55, $p < 0.01$), and lighting (54.8% inadequate; OR: 1.68, $p = 0.005$). Central government medical colleges also showed higher odds of inadequacy in several areas, including room space (73.8% inadequate; OR: 1.66, $p = 0.013$), though less pronounced than in state government medical colleges. Overall, state government

Table 3. Comparison of the Adequacy of Duty Room Amenities across Different Types of Medical Colleges in India

Duty Room Amenities	Type of Medical College (N, %)	Adequate n (%)	Inadequate n (%)	Odds Ratio	p Value
Air conditioning	Central government (377, 30.8)	230 (61.0)	147 (39.0)	0.4 (0.27–0.59)	< 0.01
	State government (697, 56.9)	160 (23.0)	537 (77.0)	2.12 (1.46–3.07)	< 0.01
	Private (150, 12.3)	58 (38.7)	92 (61.3)	Reference	-
Room space	Central government (378, 30.8)	99 (26.2)	279 (73.8)	1.66 (1.11–2.48)	0.013
	State government (697, 56.9)	136 (19.5)	561 (80.5)	2.43 (1.66–3.55)	< 0.01
	Private (151, 12.3)	56 (37.1)	95 (62.9)	Reference	-
Ventilation	Central government (377, 30.9)	112 (29.7)	265 (70.3)	1.3 (0.87–1.94)	0.192
	State government (692, 56.7)	157 (22.7)	535 (77.3)	1.88 (1.29–2.74)	< 0.01
	Private (152, 12.4)	54 (35.5)	98 (64.5)	Reference	-
Cleanliness	Central government (380, 30.9)	152 (40.0)	228 (60.0)	1.15 (0.79–1.68)	0.469
	State government (698, 56.7)	143 (20.5)	555 (79.5)	2.98 (2.06–4.31)	< 0.01
	Private (152, 12.4)	66 (43.4)	86 (56.6)	Reference	-
Pest control	Central government (375, 30.8)	92 (24.5)	283 (75.5)	1.45 (0.95–2.19)	0.081
	State government (693, 56.9)	65 (9.4)	628 (90.6)	4.55 (2.96–6.97)	< 0.01
	Private (150, 12.3)	48 (32.0)	102 (68.0)	Reference	-
Lighting	Central government (376, 30.9)	241 (64.1)	135 (35.9)	0.77 (0.53–1.14)	0.193
	State government (691, 56.8)	312 (45.2)	379 (54.8)	1.68 (1.17–2.4)	< 0.01
	Private (150, 12.3)	87 (58.0)	63 (42.0)	Reference	-

medical colleges reported the poorest adequacy across all amenities, indicating a significant disparity in facility standards compared to private institutions (Table 3).

Satisfaction with Security Measures in Healthcare Settings

The majority of the respondents expressed dissatisfaction with the existing security measures across various health institutions with regard to the availability of security personnel (28.4% dissatisfied, 24.2% very dissatisfied), surveillance and monitoring through CCTV cameras (24.2% dissatisfied, 29.6% very dissatisfied), presence of access control systems (restricted areas, use of badges and ID cards) (23.6% dissatisfied, 42.6% very dissatisfied), emergency alarms and protocols (24.2% dissatisfied, 41.5% very dissatisfied), security




Figure 4. Perception regarding Adequacy of Security Measures in Healthcare Settings in India

in areas with high-risk patients (emergencies, psychiatric wards, intensive care units) (25.4% dissatisfied, 37.2% very dissatisfied) and illumination around the working area (21.6% dissatisfied, 22.4% very dissatisfied) (Figure 4).

The alarm system for emergencies was reported to be

inadequate by 62.7% of the respondents. Around three-fourths (72.4%) of the respondents unequivocally felt that the security personnel currently stationed in these areas are not effective in handling potential threats. In comparison, 8.7% of the respondents felt that the security measures and personnel were effective, while 18.9% were not sure about the effectiveness of security.

More than 90% of health institutions (91.1%) did not have any screening mechanism or a provision for frisking to check for possession of weapons or other dangerous objects during entry to the hospital. Almost three-fourths (72.8%) of the respondents reported about the lack of a secure boundary of the hospital or health institution campus.

Variations in Satisfaction with Security Measures among Medical Colleges

The satisfaction with security services varied significantly across different types of medical colleges. Private medical colleges reported the highest satisfaction levels for all security parameters and served as the reference group. In contrast, state government medical colleges showed significantly higher dissatisfaction with the number of security personnel (63.2% dissatisfied; OR: 4.09, $p < 0.01$), monitoring/ CCTV surveillance (62.7% dissatisfied; OR: 3.78, $p < 0.01$), access control (72.5% dissatisfied; OR: 3.54, $p < 0.01$), alarms for emergencies (69.5% dissatisfied; OR: 3.52, $p < 0.01$), security in high-risk patient areas (69.4% dissatisfied; OR: 3.46, $p < 0.01$), and ambient lighting

Table 4. Comparison of Satisfaction with Security Services across Different Types of Medical Colleges in India

Satisfaction with Security Services	Type of College (N, %)	Satisfied n (%)	Dissatisfied n (%)	Odds Ratio	p Value
Number of security personnel	Central government (396, 31.1)	231 (58.3)	165 (41.7)	1.7 (1.1–2.5)	< 0.01
	State government (720, 56.5)	265 (36.8)	455 (63.2)	4.0 (2.8–5.9)	< 0.01
	Private (159, 12.5)	112 (70.4)	47 (29.6)	Reference	-
Monitoring/ CCTV surveillance	Central government (394, 31.1)	194 (49.2)	200 (50.8)	2.3 (1.5–3.4)	< 0.01
	State government (714, 56.4)	266 (37.3)	448 (62.7)	3.7 (2.6–5.4)	< 0.01
	Private (159, 12.5)	110 (69.2)	49 (30.8)	Reference	-
Access control	Central government (393, 31)	148 (37.7)	245 (62.3)	2.2 (1.5–3.2)	< 0.01
	State government (716, 56.6)	197 (27.5)	519 (72.5)	3.5 (2.4–5.0)	< 0.01
	Private (157, 12.4)	90 (57.3)	67 (42.7)	Reference	-
Alarms for emergencies	Central government (392, 31)	155 (39.5)	237 (60.5)	2.3 (1.6–3.4)	< 0.01
	State government (714, 56.5)	218 (30.5)	496 (69.5)	3.5 (2.4–5.0)	< 0.01
	Private (158, 12.5)	96 (60.8)	62 (39.2)	Reference	-
Security in areas with high-risk patients	Central government (390, 31.2)	160 (41.0)	230 (59.0)	2.1 (1.5–3.2)	< 0.01
	State government (706, 56.5)	216 (30.6)	490 (69.4)	3.4 (2.4–4.9)	< 0.01
	Private (154, 12.3)	93 (60.4)	61 (39.6)	Reference	-
Ambient lighting	Central government (395, 31.1)	232 (58.7)	163 (41.3)	1.5 (1.01–2.1)	0.04
	State government (717, 56.4)	348 (48.5)	369 (51.5)	2.2 (1.5–3.2)	< 0.01
	Private (159, 12.5)	108 (67.9)	51 (32.1)	Reference	-

(51.5% dissatisfied; OR: 2.25, $p < 0.01$). Central government medical colleges also reported increased dissatisfaction across these parameters, but to a lesser extent than state government medical colleges, with odds ratios ranging from 1.49 to 2.37. Overall, state government medical colleges exhibited the highest dissatisfaction levels, particularly for security personnel and surveillance measures as detailed in Table 4.

Incident Reporting and Emergency Response

Four-fifths (81.3%) of the respondents had been witness to violence against healthcare workers. Almost half (44.1%) of these respondents felt that the incident was not handled effectively at all and 38.3% felt that the situation was only handled partially.

With regards to action during an emergency violent situation, 80% of the respondents did not have the contact

number of the concerned security personnel on duty or did not know whom to contact when such a situation arises. In case of any security concern, 70.2% of respondents reported the lack of a clear and confidential process for reporting such concerns and 20% of respondents reported that they




Figure 5. Incident Reporting and Emergency Response (N = 1566)

were not aware whether such a process or protocol existed or not. Out of 1566 respondents, only 9.8% of respondents were aware of the hospital or health institution’s emergency protocols in case of fires, disasters, active shootings, or other violent incidents with 45.1% being “partially aware” or unsure about them. The majority of the respondents (91.6%) had not received any training regarding response to security threats and active hostile situations (Figure 5).

Qualitative Analysis

A thematic analysis of the open-ended questions identified the reasons why healthcare workers faced security threats at their place of work. It was found that agitated patients or their attendants were the most common perpetrators of threats, whereas the lack of an adequate number of security personnel and CCTV cameras further aggravated the issue. Further, unrestricted access and unauthorised entry into medical facilities compromised safety, while inadequate infrastructure such as lack of dedicated duty




Figure 6. A Tree Map of Factors Contributing to Security Concerns in Healthcare Settings in India

rooms, washrooms and ambient lighting exposed the healthcare workers to potential harm. Poor management and administrative indifference towards the lack of adequate security were also identified as reasons for feeling unsafe at work. Lastly, increasing patient load and resource constraints led to agitation among patients thus leaving the healthcare workers vulnerable to potential violence. A frequency analysis of the major themes is depicted in the tree map shown in Figure 6.

Discussion

The current study reported the perceived workplace security and concerns of healthcare professionals and analysed the security concerns in the context of gender-based violence. Perceived safety refers to the sense of security and an absence of anxiety about becoming victimised while moving through spaces¹² in contrast to actual social safety which is related to the actual crime rate¹³. Workplace security is a form of healthcare professionals’ psychological capital and a secure environment known to provide patients with more complete, comprehensive, and personalised services.^{14,15} Perception of the workplace has been found to affect job satisfaction and turnover.¹⁶

The majority of respondents did not feel safe within the hospital or healthcare institution premises during their duty hours. This perception of lack of security extended into the night shifts.

Studies from India and the Indian subcontinent have shown the prevalence of workplace violence against doctors to lie between 75% and 80%.^{17,18} The forms of threats of violence experienced by both genders are also different. Studies have reported that male doctors have a greater propensity for facing physical violence.^{19,20} The current study showed that on the whole, more females face verbal, sexual and other types of threats of violence than males but male doctors faced higher threats or instances of physical violence. Studies have attributed this difference to cultural beliefs among people against carrying out objective physical violence against women.²¹ Further exploration of the proportion of males present in workplace violence-prone areas needs to be done to explain the phenomenon better. The patriarchal dominance theory can be used to understand the rising instances of sexual violence. Also, the rise in violence in society has led to an increase in violence against doctors, particularly women, as supported by the cultural spillover concept.²⁰ It was also noted that female healthcare professionals had a lower perception of safety compared to their male counterparts. A rise in female awareness and a larger platform to speak and voice out concerns may have contributed to better identification and enlisting of security gaps. Studies have shown females to be at a higher risk of workplace violence.²² Gender dynamics, fear of retaliation and economic insecurity

have been reported as causes for women feeling unsafe at work and taking up low-paying, low-security jobs in critical areas like the emergency wards.²³ About two-thirds of the global healthcare workforce consists of women. Provisions regarding the same need to be made and the security needs of all healthcare professionals need to be accommodated.^{24,25}

The reported perceived lack of safety was found to be higher in healthcare professionals employed in state government health institutions compared to central and private health institutions. Caseload, availability of treatment modalities, and quality of security measures may be the probable causes. There have been quality differences reported between public and private hospitals with regard to resources and the quality of services.²⁶ Studies have also focused on the dearth of administrative personnel compared to healthcare professionals, thus leading to an added burden and lag in the provision of proper care and smooth workflow.²⁶ For a country like India, where the majority face the risk of catastrophic health expenditure in private hospitals,²⁷ it becomes imperative to enable public hospitals to handle the huge caseloads, especially at high-risk and critical care points like emergencies and intensive care units.^{28,29}

The study found that professionals working in government colleges and institutions were more likely to feel dissatisfied with the basic amenities like room space, ventilation, dedicated toilets, and objective security measures compared to private institutions. Among the government institutions, state government institutions and hospitals were more at risk than central government institutions highlighting the need to ramp up the same. Good duty amenities and security measures are associated with better workplace safety.³⁰ The three-tier health system in India needs to be strengthened at primary and secondary levels with adequate treatment modalities and logistics along with manpower.^{31,32} The study found doctors in primary and secondary healthcare settings to be more at risk of feeling unsafe at their workplace, wherein they are the designated first point of contact for the health system.

Interns and junior residents constituted the respondents who felt the most unsafe, while on duty. A higher duration of patient interaction and longer duty hours are the probable cause for the same. Similar studies reported resident doctors and interns being the most exposed to the risk of workplace violence, especially during night shifts.^{3,33} Long working hours put residents at a risk of burnout, which has been shown to affect productivity and efficiency. Critical areas like emergency wards demand attention and quick thinking and thus over-worked residents and interns stand at the risk and mercy of a volatile and emotional mob.^{34,35} The results from the thematic analysis in the current study put up a similar picture with the respondents feeling the most

threatened by an agitated and uncontrolled mob.

Perception of safety is important as it has been shown to affect patient management and the psycho-social well-being of the doctors. Poor patient management can perpetuate a vicious cycle of distrust and poor doctor-patient relationships, thus leading to poorer health outcomes.^{19,36} The lack of a proper and specific law protecting healthcare workers against violence often emboldens aggrieved patients or their attendants to take matters into their own hands.³⁷ Countries like Nepal,³⁸ China,³⁹ and England⁴⁰ have set a precedent.

Security and safety at the workplace are seldom treated as an area of concern for various reasons.^{41,42} Lack of funds or resources and relaxed norms could be a possible reason for the poor status of security and safety services in healthcare settings. Hence, it can be recommended that establishing reward systems, similar to the Kayakalp Award,⁴³ can not only lift morale but also provide financial incentives dedicated to improving security. Moreover, certification through mechanisms like the National Quality Assurance Standards (NQAS)⁴⁴ should be pursued to ensure high security and quality standards.

The strengths of the study include a large number of responses received from healthcare professionals working in diverse capacities and at various positions from all over India. The recent incidence of workplace violence which garnered national and international attention could have affected the responses but the objective to study this sensitive issue of safety and security required the focussed attention of the affected party. Open-ended questions have captured qualitative aspects of information. Therefore, the high sensitivity of the assessing questionnaire should be accepted as beneficial for the study.

Limitations

The use of a non-probability sampling technique may have compromised the representativeness of the study, but we have tried to get a bigger sample size to avoid recruitment bias. The usage of Google Forms as a form of data collection may have led to selection bias with more technology-friendly respondents opting to fill out the form. There may have been some form of recall bias due to the unspecified duration of experiencing workplace violence.

Recommendations

To enhance security measures in medical and educational institutions, several key recommendations are proposed. Firstly, improving the conditions of duty rooms with all necessary amenities, particularly in state and central government institutions, will ensure adequate safety and rest for the healthcare professional. Locally, institute

administration should ensure that an appropriate number of security guards are maintained working in tandem with local police to handle threats and increase their effectiveness. Implementing an effective local system of systems (SOS) will provide immediate response options in emergencies while securing the perimeter of the healthcare setting will minimise risk and improve access control. The security in areas with high-risk patients such as the emergency wards, psychiatry wards, and intensive care units needs to level up from the current status. Security at night can be upgraded by identifying blind spots and installing lights and CCTV cameras. A transport vehicle can be used to travel within the campus at night. Healthcare institutions should establish clear protocols for handling violence, provide accessible emergency contacts, and implement a confidential reporting system for security concerns. Regular training and drills on emergency response and hostile situations should be made essential, while regular awareness campaigns and security audits are essential to evaluate and enhance these measures effectively. The administration at state government medical colleges should work for a comprehensive upheaval of security measures as they were found lagging in several parameters. Strengthening of security measures is also required at the peripheral levels such as community health centres, primary health centres, and district hospitals. Special communication workshops can be held for undergraduate students to improve communication and foster a good doctor-patient relationship.^{45,46} Inter-sectoral coordination among national agencies such as the National Health Systems Resource Centre (NHSRC),⁴⁷ a premier think tank for the Ministry of Health and Family Welfare (MoHFW), National Medical Commission (NMC),⁴⁸ National Institute of Occupational Health (NIOH),⁴⁹ and central and state law agencies including the local police will help build and sustain effective mechanisms to ensure workplace safety for healthcare workers. Security considerations should be integrated into the development plans of new medical colleges and healthcare units, ensuring that safety is prioritised from the outset. Policy-level changes are also needed to make security a priority, supported by special legislation at central and state levels, to provide a standardised legal framework for protection across all institutions.

Conclusion

The findings of this study reveal that more than half of the health workforce perceive their workplace as unsafe. The conditions of the duty rooms, especially in the state and central government medical colleges, are grossly inadequate. Healthcare professionals working in public hospitals reported feeling more unsafe compared to those in the private sector. Improving workplace security in healthcare settings requires a comprehensive, need-based approach with strict laws to safeguard healthcare personnel. Assurance of workplace safety and security shall help in improving doctor-patient

relationships, building a healthier workforce, and providing a better quality of healthcare services to the community.

Acknowledgement

We are grateful to Dr Jugal Kishore for his guidance. We are also thankful to all participants for their prompt consents, responses and suggestions.

Source of Funding: None

Conflict of Interest: None

References

1. Legislative Department India [Internet]. Constitution of India; [cited 2024 Sep 14]. Available from: <https://legislative.gov.in/constitution-of-india/>
2. World Health Organization [Internet]. Health workforce; [cited 2024 Aug 26]. Available from: <https://www.who.int/india/health-topics/health-workforce>
3. Kumar M, Verma M, Das T, Pardeshi G, Kishore J, Padmanandan A. A study of workplace violence experienced by doctors and associated risk factors in a tertiary care hospital of South Delhi, India. *J Clin Diagn Res.* 2016 Nov;10(11):LC06-10. [PubMed] [Google Scholar]
4. Bhatia N, Kishore J, Anand T, Jiloha RC. Occupational stress amongst nurses from two tertiary care hospitals in Delhi. *Australas Med J.* 2010 Jan 1;3(11):731-8. [Google Scholar]
5. Gupta M, Kishore J, Kohli C. Rural posting for medical graduates: perception, acceptance and plausibility. *Int J Healthc Educ Med Inform.* 2017;4(1):3-8. [Google Scholar]
6. Berlanda S, Pedrazza M, Fraizzoli M, de Cordova F. Addressing risks of violence against healthcare staff in emergency departments: the effects of job satisfaction and attachment style. *Biomed Res Int.* 2019;2019:5430870. [PubMed] [Google Scholar]
7. Fida R, Tramontano C, Paciello M, Guglielmetti C, Gilardi S, Probst TM, Barbaranelli C. 'First, do no harm': the role of negative emotions and moral disengagement in understanding the relationship between workplace aggression and misbehavior. *Front Psychol* [Internet]. 2018 May 11 [cited 2024 Aug 26];9:671. Available from: <https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2018.00671/full> [PubMed] [Google Scholar]
8. Sachdeva S, Jamshed N, Aggarwal P, Kashyap SR. Perception of workplace violence in the emergency department. *J Emerg Trauma Shock.* 2019;12(3):179-84. [PubMed] [Google Scholar]
9. Hamdan M, Abu Hamra A. Workplace violence towards workers in the emergency departments of Palestinian hospitals: a cross-sectional study. *Hum Resour Health.* 2015 May 7;13:28. [PubMed] [Google Scholar]
10. O'Brien CJ, van Zundert AA, Barach PR. The growing

- burden of workplace violence against healthcare workers: trends in prevalence, risk factors, consequences, and prevention – a narrative review. *EClinicalMedicine*. 2024;72:102641. [PubMed] [Google Scholar]
11. Ministry of Health & Family Welfare [Internet]. ServicePlus- Suggestions to NTF on safety, working conditions and well-being of medical professionals and other cognate matters; [cited 2024 Sep 15]. Available from: https://serviceonline.gov.in/renderApplicationForm.do;jsessionid=6038B49978955069F72F2E3454D9D320?serviceId=29870002&UUIID=aaa66f73-705e-4f56-8e16-0b9009eb0bf7&directService=true&templd=14682&serviceNameToDisplay=Suggestions+to+NTF+on+safety%2C+working+conditions+and+well-being+of+medical+professionals+and+other+cognate+matters&grievDefined=0&srviceLinkRequired=No&userLoggedIn=N&source=CTZN&OWASP_CSRFTOKEN=DIJD-YZ8W-H68J-Z5EX-QHPD-UXN3-O76Q-3B7P
 12. Boomsma C, Steg L. Feeling safe in the dark: examining the effect of entrapment, lighting levels, and gender on feelings of safety and lighting policy acceptability. *Environ Behav*. 2014 Feb 1;46(2):193-212. [Google Scholar]
 13. Park AJ, Calvert TW, Brantingham PL, Brantingham PJ. The use of virtual and mixed reality environments for urban behavioural studies. *PsychNol J* [Internet]. 2008 Jan 1 [cited 2024 Aug 26];6(2):119-30. Available from: https://www.academia.edu/14875478/The_Use_of_Virtual_and_Mixed_Reality_Environments_for_Urban_Behavioural_Studies [Google Scholar]
 14. Xu Y, Zhan N, Zhang D, Xie Z, Li G, Huang H. The effect of sense of security on job performance of medical staff: the mediating effect of psychological capital. *Front Psychol*. 2024 Apr 4;15:1347783. [PubMed] [Google Scholar]
 15. World Health Organization [Internet]. Keep health workers safe to keep patients safe: WHO; 2020 [cited 2024 Sep 14]. Available from: <https://www.who.int/news/item/17-09-2020-keep-health-workers-safe-to-keep-patients-safe-who>
 16. Ahamed F, Kaur A, Sengupta P, Ghosh T. Perception of safety from workplace violence affects job satisfaction among doctors practicing modern medicine in India: a nationwide survey. *J Family Med Prim Care*. 2021 Jun;10(6):2252-8. [PubMed] [Google Scholar]
 17. Rony MK, Sharmi PD, Parvin MR, Kayesh I, Alamgir HM. Prevalence and risk factors of workplace violence against healthcare workers in Bangladesh and its consequences: a nationwide cross-sectional study. *Inform Med Unlocked*. 2023 Aug 28;41:101335. [Google Scholar]
 18. Debnath A, Alam M, Goyal M, Khokhar A, Lukhmana S. The prevalence of violence against resident doctors and its subsequent effects on patient management in a tertiary care hospital in Delhi, India. *Cureus*. 2023 May 17;15(5):e39116. [PubMed] [Google Scholar]
 19. Singh A, Ranjan P, Agrawal R, Kaur T, Upadhyay AD, Nayer J, Chakrawarty B, Sarkar S, Joshi M, Kaur TP, Mohan A, Chakrawarty A, Kumar KR. Workplace violence in healthcare settings: a cross-sectional survey among healthcare workers of North India. *Indian J Occup Environ Med*. 2023;27(4):303-9. [PubMed] [Google Scholar]
 20. Nainar V. Understanding the rise of sexual violence in India [Internet]. Policy Press; 2022 Feb 1 [cited 2024 Aug 26]. p. 9-27. Available from: <https://bristoluniversitypressdigital.com/view/journals/jgbv/6/1/article-p9.xml> [Google Scholar]
 21. Sun L, Zhang W, Qi F, Wang Y. Gender differences for the prevalence and risk factors of workplace violence among healthcare professionals in Shandong, China. *Front Public Health*. 2022;10:873936. [PubMed] [Google Scholar]
 22. Parodi JB, Burgos LM, Garcia-Zamora S, Liblik K, Pulido L, Gupta S, Saldarriaga C, Puente-Barragan AC, Morejon-Barragan P, Alexanderson-Rosas E, Sosa-Liprandi A, Botto F, Sosa-Liprandi MI, Lopez-Santi R, Vazquez G, Gulati M, Branachuk A. Gender differences in workplace violence against physicians and nurses in Latin America: a survey from the Interamerican Society of Cardiology. *Public Health*. 2023 Dec 1;225:127-32. [PubMed] [Google Scholar]
 23. Lancet T. The structural roots of violence against female health workers. *Lancet*. 2024 Sep 7;404(10456):907. [PubMed] [Google Scholar]
 24. Jayadevan R, Augustine D, Anithadevi TS, Ramachandran R, Benaven J. Safety during night duty: survey of 3885 doctors across India. *medRxiv* [Preprint]; 2024 [cited 2024 Sep 12]. p. 2024.09.03.24312775. Available from: <https://www.medrxiv.org/content/10.1101/2024.09.03.24312775v1> [Google Scholar]
 25. World Health Organization [Internet]. Value gender and equity in the global health workforce; [cited 2024 Sep 12]. Available from: <https://www.who.int/activities/value-gender-and-equity-in-the-global-health-workforce>
 26. Tiwari RV, Sharma SK, Sahoo SR, Velthuru SK, Basavarajaiah JM, Kazi M, Dixit H. Comparative evaluation of quality management practices in the public and private hospitals in Visakhapatnam district: an original research. *J Pharm Bioallied Sci*. 2024 Feb;16(Suppl 1):S592-7. [PubMed] [Google Scholar]
 27. Poverty and Inequality Platform [Internet]. Country profile: India; [cited 2024 Sep 12]. Available from:

- <https://pip.worldbank.org/country-profiles/IND>
28. Anand T, Grover S, Kumar R, Kumar M, Ingle GK. Workplace violence against resident doctors in a tertiary care hospital in Delhi. *Natl Med J India*. 2016 Nov 1;29(6):344. [PubMed] [Google Scholar]
 29. Kharche J, Jacob R, Patel S. To study the violence experienced by postgraduate resident doctors and interns in a tertiary care centre in Aurangabad. *Asian J Med Health*. 2022;20(6):25-9. [Google Scholar]
 30. Morken T, Johansen IH. Safety measures to prevent workplace violence in emergency primary care centres—a cross-sectional study. *BMC Health Serv Res*. 2013 Oct 3;13:384. [PubMed] [Google Scholar]
 31. Khan A, Yattoo G, Mir MS. Why our tertiary care hospital emergencies are overcrowded? Faulty referrals or anything else: a study from a teaching hospital of North India. *Int J Adv Res Innov Ideas Educ*. 2021 Feb 17;7(1):345-50.
 32. Ghia C, Rambhad G. Implementation of equity and access in Indian healthcare: current scenario and way forward. *J Mark Access Health Policy*. 2023 Mar 26;11(1):2194507. [PubMed] [Google Scholar]
 33. Nayanar BS, Fareed N, Battur H, Praveena J. A study on nature of violence against doctors in tertiary care centers in Karnataka, India: a cross-sectional study. *Indian J Community Med*. 2024;49(3):472. [PubMed] [Google Scholar]
 34. Amer SA, Elotla SF, Ameen AE, Shah J, Fouad AM. Occupational burnout and productivity loss: a cross-sectional study among academic university staff. *Front Public Health*. 2022 Apr 25;10:861674. [PubMed] [Google Scholar]
 35. Kovacs R, Lagarde M. Does high workload reduce the quality of healthcare? Evidence from rural Senegal. *J Health Econ*. 2022 Mar;82:102600. [PubMed] [Google Scholar]
 36. Kaur A, Ahamed F, Sengupta P, Majhi J, Ghosh T. Pattern of workplace violence against doctors practising modern medicine and the subsequent impact on patient care, in India. *PLoS One*. 2020 Sep 18;15(9):e0239193. [PubMed] [Google Scholar]
 37. Nair A, Zadey S. Ending violence against healthcare workers in India: a bill for a billion. *Lancet Reg Health Southeast Asia* [Internet]. 2022 Nov [cited 2024 Sep 12];6:100064. Available from: [https://www.thelancet.com/journals/lansea/article/PIIS2772-3682\(22\)00080-4/fulltext](https://www.thelancet.com/journals/lansea/article/PIIS2772-3682(22)00080-4/fulltext) [PubMed] [Google Scholar]
 38. Kharel S. A historic ordinance against violence to health workers of Nepal. *Lancet Reg Health Southeast Asia* [Internet]. 2022 Aug [cited 2024 Sep 12];3:100037. Available from: [https://www.thelancet.com/journals/lansea/article/PIIS2772-3682\(22\)00047-6/fulltext](https://www.thelancet.com/journals/lansea/article/PIIS2772-3682(22)00047-6/fulltext) [PubMed] [Google Scholar]
 39. Lu S, Ren S, Xu Y, Lai J, Hu J, Lu J, Huang M, Ma X, Chen J, Hu S. China legislates against violence to medical workers. *Lancet Psychiatry*. 2020 Mar;7(3):e9. [PubMed] [Google Scholar]
 40. Legislation.gov.uk [Internet]. Medical Act 1983; [cited 2024 Sep 12]. Available from: <https://www.legislation.gov.uk/ukpga/1983/54/contents>
 41. Huang YH, Leamon TB, Courtney TK, Chen PY, DeArmond S. Corporate financial decision-makers' perceptions of workplace safety. *Accid Anal Prev*. 2007;39(4):767-75. [PubMed] [Google Scholar]
 42. Ahasan R. Legacy of implementing industrial health and safety in developing countries. *J Physiol Anthropol Appl Human Sci*. 2001;20(6):311-9. [PubMed] [Google Scholar]
 43. National Health Systems Resource Centre [Internet]. Kayakalp and Swachh Swasth Sarvatra; [cited 2024 Sep 8]. Available from: <https://qps.nhsrindia.org/kayakalp-swachh-swasth-sarvatra>
 44. National Health Systems Resource Centre [Internet]. National Quality Assurance Standards; [cited 2024 Sep 8]. Available from: <https://qps.nhsrindia.org/national-quality-assurance-standards>
 45. Ha JF, Longnecker N. Doctor-patient communication: a review. *Ochsner J*. 2010;10(1):38-43. [PubMed] [Google Scholar]
 46. Sethi S, Pardeshi G, Kishore J. Communication skills for interns: evolution and assessment of a training workshop in a medical college of New Delhi. *Int J Healthc Educ Med Inform*. 2019;6(2):17-22. [Google Scholar]
 47. National Health Systems Resource Centre [Internet]. About us; [cited 2024 Sep 8]. Available from: <https://nhsrindia.org/about-us>
 48. National Medical Commission [Internet]. About NMC; [cited 2024 Sep 8]. Available from: <https://www.nmc.org.in/about-nmc/introduction/>
 49. National Institute of Occupational Health [Internet]. About NIOH Ahmedabad; [cited 2024 Sep 8]. Available from: <https://nioh.org/nioh-ahmedabad/>